



Online Survey on the Acceptance of Latent Tuberculosis Testing and Isoniazid Preventive Therapy among Healthcare Workers in Nigeria

Solomon Obinna Obidigbo ^a, Fatima Ibrahim Makintami ^b,
Yusuf Madaki Lekko ^c, Mohammed Usman Sajo ^d
and Francis Eneche Ejeh ^{a,d*}

^a Department of Public Health, Maiduguri Study Center, National Open University of Nigeria, Nigeria.

^b Nigeria Agriculture Quarantine Services (NAQS), Nigeria.

^c Department of Veterinary Medicine, University of Maiduguri, Nigeria.

^d Department of Veterinary Microbiology, University of Maiduguri, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author FEE conceptualized. Authors FEE, MUS and SO did the methodology, did formal analysis, did the investigation, did the resources.

FEE did the software. Authors FEE and MUS did the data curation. Author FEE wrote original draft. Authors FEE, SO, MUS and YML reviewed and edited the manuscript. FEE supervised the study. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpr/2024/v13i5308>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/122966>

Original Research Article

Received: 05/07/2024

Accepted: 10/09/2024

Published: 13/09/2024

*Corresponding author: Email: enenche francis@unimaid.edu.ng, enenche francis@yahoo.com;

Cite as: Obidigbo, Solomon Obinna, Fatima Ibrahim Makintami, Yusuf Madaki Lekko, Mohammed Usman Sajo, and Francis Eneche Ejeh. 2024. "Online Survey on the Acceptance of Latent Tuberculosis Testing and Isoniazid Preventive Therapy Among Healthcare Workers in Nigeria". *International Journal of Pathogen Research* 13 (5):41-53. <https://doi.org/10.9734/ijpr/2024/v13i5308>.

ABSTRACT

Background: Healthcare workers in Nigeria are exposed to myriads of occupational risks including workplace acquired infections such as tuberculosis.

Methods: This study was a cross-sectional survey and used google form to administer the questionnaire via social media networks. Descriptive statistical methods were used to summarize data on socio-demographic characteristics and responses to questions, Microsoft excel and SPSS version 20.0 for statistical analysis of data summarized as categorical variables, frequency (n) and percentages (%).

Results: About six out of ten of the participants believed that they could be severely ill if they got tuberculosis. Majority of respondents 413 (89.2%) reported acceptance of latent TB testing while 87.7% agreed to take Isoniazid treatment if they test positive for LTBI. The occupational risk group was found to be less likely to accept LTBI testing. Also, respondents from the Northeast (OR = 4.225; 95% CI = 1.216-14.688; p value = 0.023) and southeast (OR = 4.494; 95% CI = 1.382-14.609; p value = 0.012) were four times more likely to accept isoniazid preventive treatment respectively. Participants who held that tuberculosis is a serious disease were more likely to accept isoniazid preventive treatment (OR = 2.220; 95% CI = 1.100-4.480 p value = 0.026). The majority of the HCW (56%) in the study never get tested for LTBI.

Conclusion: the majority of the healthcare workers in this study showed that TB is a dangerous disease and agreed to get regular testing and isoniazid therapy to prevent tuberculosis.

Keywords: Isoniazid preventive therapy; latent tuberculosis; vaccination; BCG vaccine; vaccine acceptance.

1. BACKGROUND

The continuous rise in global tuberculosis incidence, emergence of multi-drug and extensive drug-resistant tuberculosis and the increased occupational risk being posed to healthcare workers (HCWs) suggests that current global TB control and prevention strategies were inadequate. The World Health Organization (WHO) proposed the directly observed treatment short course (DOTS) strategy for effective TB control. The key elements of the strategy are political commitment to support TB treatment, detection of TB by sputum microscopy, molecular detection of tuberculosis and rifampicin resistant tuberculosis, multi-drug-resistant tuberculosis, direct observation of short course therapy, regular supply of medicines and reporting of programme performance and treatment outcomes. To eliminate tuberculosis in Nigeria the DOTs programme was launched in 1993, since then there has been a slow increase in tuberculosis case detection, contact tracing and treatment. However, the emergence of COVID-19 pandemic has led to a reverse in the gain made against tuberculosis in Nigeria and worldwide [1].

Latent TB infection (LTBI) refers to the state where one's immune system is continuously activated by the TB bacilli, but they are clinically

not sick [2]. Prevention of the development of active TB in healthcare workers is important since they are constantly at risk of transmitting tuberculosis to their patients and acquiring it from them. Also, HCWs who have latent tuberculosis coinfecting with HIV have a 10% yearly risk of developing active TB as compared to 10% lifetime risk in the HIV negative population [3]. HIV Virus infection was reported to be associated with impaired TB-specific CD4+ T-cell function and proliferative capacity [4]. Functionally impaired CD4+ T cells may be unable to control TB replication [5], particularly when this is combined with an increased level of an anti-inflammatory cytokine IL-10 in the serum [5]. TB and malaria weakened immune responses to TB [6]. Also, HCV has an inhibitory effect on the immune response against TB [7]. Screening of HCWs for latent TB and the provision of preventive treatment are components of the first pillar of the World Health Organization's (WHO) End TB Strategy [8]. It is hoped that by the year 2030 the World Health Organization and its partner will end tuberculosis epidemic globally [1].

Occupational TB is expected to be high among healthcare workers in Nigeria as the countries is experiencing shortage in health workforce and facilities which is further exacerbated by the continuous increase in population growth and dwindling resources. And putting undue pressure

on available HCWs. Therefore, understanding the preventive measure adapted by healthcare workers and predisposing risk factors of active TB disease among healthcare workers in Nigeria is important in managing the disease in the establishment. Such information may lead to specific and targeted responses to disease control.

The priority groups for LTBI testing and treatment have a higher prevalence of LTBI and risk of advancement to active TB than the general population. Compared to the LTBI burden in the general population worldwide (23.0%) [9], the prevalence of LTBI among healthcare workers in low/middle-income countries was stated to be 49% with the tuberculin skin test and 39% with the interferon γ release assay in a systematic review [10]. Moreover, individuals with immunocompromised status have higher risk of progression to active tuberculosis.

Worldwide about 10.6 million people developed TB in the year 2022, and Nigeria accounted for about 4.5% of people who developed TB in 2023 [1]. The prevalence of latent tuberculosis among healthcare workers in Nigeria was high in a study conducted in Akwa-Ibom state of Nigeria, [11] reported a prevalence of LTBI was 24.8% and 45.8% as assessed by QFT-IT and TST respectively. Healthcare workers are the most important frontliners in the fight against tuberculosis. Yet, this category of the population is often neglected with regards to tuberculosis control and prevention. Hence this study aimed to evaluate the factors associated with latent tuberculosis testing and acceptance of isoniazid preventive therapy among healthcare workers in Nigeria.

2. MATERIALS AND METHODS

2.1 Study Design

We commenced a cross-sectional, web-based anonymous survey using an online questionnaire. The researchers used the social network platforms of Facebook, Instagram, and WhatsApp to disseminate and advertise the survey link to the healthcare workers and health support staff. The inclusion criteria were that the respondents were healthcare workers, healthcare support staff who are Nigerian residents, and were 25 years old and above.

2.2 Study Population

This study was conducted at the six geopolitical zones in Nigeria. The study was reviewed and

approved by the committee of the research supervisors at the National Open University of Nigeria (NOUN). Informed consent was implied following the completion and submission of the questionnaire.

Participants were twenty-five years and above, adults, health workers and residents in Nigeria. Healthcare workers (HCWs) in this study include medical doctors, veterinary doctors, nurses, pharmacist, medical laboratory scientist, radiologist, dentist, public health officers and others. While Healthcare support staff (HCSS) include genitors, drivers, electricians, engineers, security, health record and research, ward attendants, etc. Google form was used to communicate the questionnaire web link to the target participants. Social media platforms such as WhatsApp, Twitter, Facebook, and Instagram were used to reach participants. A brief explanation was included in the introduction of the questionnaire. The submitted answered questionnaire was considered consent to take part in the study. The questionnaire was pre-evaluated for consistency and error in accordance with [12].

2.3 The Measures

The survey instrument used in this study was based on the tool designed by Municipal Public Health Service Rotterdam-Rijnmond (GGD) together with the National Institute for Public Health and the Environment (RIVM) in the Netherlands [13] with minimal modification. The questionnaire consisted of four (3) parts:

2.3.1 Part one (1): Demographic characteristics of respondents

Participants were asked to report their age, sex, state of residence, educational level, marital status, and profession. The state of residents was later grouped into geopolitical zones. Nigeria is categorized into six geopolitical zone viz North central, Northeast, Northwest, Southeast, South-south and Southwest.

2.3.2 Part two (2): Perception of susceptibility to tuberculosis

This category included five (5) questions on the risk of acquiring tuberculosis The risk perception questions consisted of Likert questions (1, certainly not 5, most certainly). This was later converted to dichotomous questions (certainly not, most certainly); a most certainly answer was

scored 1 point, while a certainly not answer was scored 0 points. The total mean perception of susceptibility score ranged from zero to five; the mean knowledge score of 0-3.5 ($\leq 69\%$) was considered certainly not susceptible and a mean score of 3.6-5.0 (70% and above) represent most certain of the susceptibility to tuberculosis.

2.3.3 Part three (3) Perceived severity to tuberculosis

The participant's perception of the severity of tuberculosis was measured by three items (e.g., "I am afraid of even thinking about getting ill with tuberculosis"; "I could be severely ill if I got tuberculosis"; How likely do you think it is that a person who falls ill with tuberculosis dies because of the disease?"). The question consisted of a 5-point Likert scale (1, Not at all serious 5, very serious). with a higher score indicating a higher level of perceived severity. The Likert scale was converted to a dichotomous scale for the purpose of calculation mean score and for ease of analysis. (Not at all serious, not serious, and not serious-slightly serious were assigned zero; that is Not at all serious while serious and very serious were scored 1; that is very serious).

2.3.4 Acceptance of screening test for latent tuberculosis and isoniazid

This item consisted of two questions (Would you accept testing for latent tuberculosis if this was advised by the government or medical practitioners? and would you accept isoniazid treatment if tested positive for latent tuberculosis?)

2.4 Statistical Analyses

Descriptive statistical methods were used to summarize data on socio-demographic characteristics and responses to questions regarding respondents' perception of the seriousness, susceptibility, and the benefit of tuberculosis immunization. And the intention to accept testing for latent tuberculosis and isoniazid treatment among healthcare workers in Nigeria. Data were entered in Microsoft Excel and later imported into SPSS version.20.0 for statistical analysis. Data were summarized as categorical variables' frequencies (n) and percentages (%). Furthermore, we presented the data in tables and figures. Chi-square was used to determine the level of difference among the

categorical variables. A binary logistic regression analysis was applied and expressed as odds ratio (OR) and 95% confidence interval (CI) to determine the predictors of the willingness to accept latent tuberculosis testing and acceptance of isoniazid preventive treatment. A p-value of less than 0.05 was considered significant in all tests.

3. RESULTS

3.1 Demographic Characteristics of the Study Population

The survey was disseminated to healthcare workers in Nigeria from April to August 2022, and a total of 463 healthcare workers completed and submitted the advertisement online. The survey received responses from participants of diverse demographic characters. Table 1 showed that the study participants had huge representation of 25-35 years old 328 (70.8%), male 327 (70.6%) university first degree 272 (58.7%) and healthcare workers 392 (84.7%) only a small portion reported high risk perceptions 241(5.2%) to tuberculosis infections.

3.2 Perception of Susceptibility

Table 2 shows the participants responses to questions regarding their perceived susceptibility to tuberculosis. Most of respondents held that they are at risk of getting tuberculosis because of their lifestyle and underlying illness were 62 (14.9%) respectively. The risk of getting tuberculosis because of family history of TB was 62 (13.4%). The occupational risk of getting tuberculosis was higher 282 (60.9%).

3.3 Perception of Severity of Tuberculosis among Healthcare Workers in Nigeria

Concerning the fear of getting tuberculosis (68.3%) of the respondents held that they were not afraid of thinking about getting ill with tuberculosis but majority 278 (60%) of the respondents believed that they could be severely ill if they got tuberculosis. However, 136 (29.4%) believed that a person who fall ill with tuberculosis could die (Table 2). Table 3 compared the perception of seriousness of tuberculosis and demographic characteristics among healthcare workers in Nigeria.

Table 1. Demographic Characteristics of the study participants

Variables		Frequency	Percentage (%)
Age group	25-35	328	70.8
	36-46	109	23.5
	47-57	22	4.8
	58 and above	4	0.9
Marital Status	Single	191	41.3
	Married	272	58.7
Gender	Male	327	70.6
	Female	136	29.4
Religion	Christianity	311	67.2
	Islam	152	32.8
Geopolitical Zone	Northcentral	73	15.6
	Northeast	213	46.0
	Northwest	25	5.4
	Southeast	111	24.0
	South-south	21	4.5
	Southwest	20	4.3
Educational Level	Secondary	26	5.6
	Diploma	28	6.1
	First degree	272	58.7
	Postgraduate degree	137	29.6
Occupation	Healthcare worker	392	84.7
	Healthcare support staff	71	15.3
Risk perception	Low risk perception	439	94.8
	High risk perception	24	5.2
Total		463	100.0




Fig. 1. Perception of susceptibility to tuberculosis among the respondents

Table 2. Perceived susceptibility, severity, benefit of tuberculosis vaccine and Acceptance of tuberculosis testing and treatment

Perceived susceptibility to tuberculosis		
Question	Responses (%)	
	Certainly not (%)	Most certainly (%)
I am at high risk of tuberculosis because of my family medical history	401 (86.6)	62 (13.4)
I am at high risk of tuberculosis because my own underlying medical conditions	394 (85.1)	69 (14.9)
I am at high risk of tuberculosis because of my occupational history	181 (39.1)	282 (60.9)
I am at high risk of tuberculosis because of my lifestyle	394 (85.1)	69 (14.9)
I think I will get tuberculosis soon	449 (97.0)	14 (3.0)
Perceived severity to tuberculosis		
	Not at all serious	Very serious
I am afraid of even thinking about getting ill with tuberculosis	316 (68.3)	149 (31.7)
I could be severely ill if I get tuberculosis	185 (40.0)	278 (60.0)
	Not likely	Most likely
How likely do you think it is that a person who falls ill with tuberculosis dies because of the disease?	327 (70.6)	136 (29.4)
Acceptance of screening test for latent tuberculosis and isoniazid treatment		
	No	Yes
Will accept testing for latent tuberculosis if this was advised by the government or medical practitioners	50 (10.8)	413 (89.2)
Will accept isoniazid treatment if tested positive for latent tuberculosis	66 (14.3)	397 (85.7)

Concerning the perception of severity of tuberculosis among healthcare workers in Nigeria, 363(68.3%) held that they were not afraid of getting ill with tuberculosis. There were significant ($p<0.05$) differences among age group regarding the fear of getting TB. The age group 47 and above reported 100% not afraid of getting TB while 214(65.2%) of the age group 25–35-year-old were afraid of getting TB.

There was statistically significant difference ($p<0.05$) among age group, marital status and religion regarding respondent who reported they could be severely ill if they get TB. The age group 25-35 reported high (64.6%) severity if they get TB.

Respondents that had never married (single) reported 1.25 (65.4%) that they could be severely ill if they get TB. Also, 201(64.6%) of the respondents in the study reported that they could be severely ill if they get TB. The perception that a person infected with TB could die was significantly ($p<0.05$) different among gender, age group and geopolitical zones. Being female gender 49(36.0%) age group fifty-nine

and above (50.0%) and respondents from the south-south geopolitical zones reported higher perception that TB could lead to death. (Table 3).

3.4 Factors associated with Acceptance of Latent tuberculosis testing by healthcare Workers in Nigeria

Four hundred and thirteen representing 89.2% of the respondents reported a definite acceptance of latent TB testing. The age group 36-46 years old (OR=14.613, 95% ci 1.320-161.833, $P=0.029$) were more likely to accept latent TB testing compared to 59 years old and above. Respondents who tested regularly for latent TB were reported to more likely to accept latent TB testing. (OR=4.062, 56ci=1.126-14653, $P=0.032$). (Table 4) participants who reported that they were at high risk of getting TB because of their occupations were less likely to accept latent TB testing (OR = 0.223 95% ci =0.106-0.471, $p=0.000$). Fig. 2 shows that a total of 259 (56.0%) responded yes to never getting tested for latent TB.

Table 3. Perception of Severity of Tuberculosis among Healthcare Workers in Nigeria

Variables	I am afraid of getting TB			I could be severely ill if I get TB			Persons infected with TB could die		
	Yes (%)	No (%)	χ^2 (p value)	Yes (%)	No (%)	χ^2 (p value)	Yes (%)	No (%)	χ^2 (p value)
Gender									
Male	106 (32.6)	221 (67.6)	0.228 (0.633)	198 (60.6)	129 (39.4)	0.911 (0.730)	87 (26.6)	240 (73.4)	4.112 (0.043)
Female	41 (30.1)	95 (69.9)		80 (58.8)	56 (41.2)		49 (36.0)	87 (64.0)	
Age group									
25-35	114 (34.8)	214 (65.2)	13.573 (0.004)	212 (64.6)	116 (35.4)	10.581 (0.014)	114 (34.8)	214 (65.2)	21.458 (0.000)
36-46	33 (30.3)	76 (69.7)		55 (50.5)	54 (49.5)		13 (11.9)	96 (88.1)	
47-57	0 (0.0)	22 (100.0)		9 (40.9)	13 (59.1)		7 (31.8)	15 (68.2)	
≥58	0 (0.0)	4 (100.0)		2 (50.0)	2 (50.0)		2 (50.0)	2 (50.0)	
Marital status									
Single	59 (30.9)	132 (69.1)	0.111 (0.739)	125 (65.4)	66 (34.6)	3.954 (0.047)	59 (30.9)	132 (69.1)	0.360 (0.548)
Married	88 (32.4)	184 (67.6)		153 (56.2)	119 (43.8)		77 (28.3)	195 (71.7)	
Religion									
Christianity	98 (31.5)	213 (68.5)	0.025 (0.875)	201 (64.6)	110 (35.4)	8.308 (0.004)	96 (30.9)	215 (69.1)	1.020 (0.313)
Islam	49 (32.2)	103 (67.8)		77 (50.7)	75 (49.3)		40 (26.3)	215 (69.1)	
Geographical zone									
Northcentral	24 (32.9)	49 (67.1)	7.667 (0.176)	48 (65.8)	25 (34.2)	4.680 (0.456)	20 (27.4)	53 (72.6)	12.218 (0.032)
Northeast	64 (30.0)	149 (70.0)		126 (59.2)	87 (40.8)		65 (30.5)	148 (69.5)	
Northwest	11 (44.0)	15 (56.0)		11 (44.0)	14 (56.0)		11 (44.0)	14 (56.0)	
Southeast	33 (29.7)	78 (70.3)		70 (63.1)	41 (36.9)		23 (20.7)	88 (79.3)	
South-south	11 (52.4)	10 (47.6)		11 (52.4)	10 (47.6)		11 (52.4)	10 (47.6)	
Southwest	4 (20.0)	16 (80.0)		12 (60.0)	8 (40.0)		6 (30.0)	14 (70.0)	
Education									
Secondary certificate	6 (23.1)	20 (76.9)	4.032 (0.258)	16 (61.5)	10 (38.5)	4.501 (0.212)	4 (15.4)	22 (84.6)	3.104 (0.376)
Diploma	8 (28.6)	20 (71.4)		12 (42.9)	16 (57.1)		10 (35.7)	18 (64.3)	
First degree	96 (35.3)	176 (64.7)		162 (59.6)	110 (40.4)		80 (29.4)	192 (70.6)	
Postgraduate	37 (27.0)	100 (73.0)		88 (64.2)	49 (35.8)		42 (30.7)	95 (69.3)	
Occupation									
Healthcare worker	129 (32.9)	263 (67.1)	1.584 (0.208)	234 (59.7)	158 (40.3)	0.130 (0.818)	122 (31.1)	270 (68.9)	3.768 (0.052)

3.5 Demographic Factors Associated with the Acceptance of Isoniazid Treatment by Healthcare Workers in Nigeria

The proportion of respondents who agreed to take isoniazid treatment whenever they are tested and return positive for latent TB was 87.7%. Respondents from the Northeast (OR, 4.225; 95%CL =1.216-14.688, P=0.023) and Southeast (OR= 4.494;95%CL =1.382-14.609 P=0.012) were about four times more likely to commence treatment with isoniazid. Educational attainment was not a significant (p<0.05) prediction of the willingness to accept isoniazid treatment by healthcare workers.

Occupational risk for getting tuberculosis was less likely to be associated with the acceptance of isoniazid treatment (OR 0.357 95% cl =0.188-

0.680, p=0.002). However, respondents who perceived Tuberculosis to be a serious disease (OR=2.220,95%cl =1.100-4.480 p=0.026) and the belief that tuberculosis vaccination prevents against infections with tuberculosis (OR=3.77;95%cl =1.230-11.605, p=0.020) were found to have the highest significant odds of the willingness to accept isoniazid preventive treatment. (Table 4).

The participants had high perceptions of the benefits of tuberculosis immunization. The majority agree that immunization prevent tuberculosis 366 (73.2%) ($\chi^2 =12.458$; p = 0.006). There was significant difference regarding group and the prevention of tuberculosis by Immunization. The age group 25-35 years old 255(77.7%) had higher proportion that agreed that immunization could prevent Tuberculosis.

Table 4. Demographic factors associated with the acceptance of latent tuberculosis testing among healthcare workers in Nigeria

Variable	OR	95% C. I	P-value
Age group			
25-35	7.438	0.702-78.785	0.096
36-46	14.613	1.320-161.833	0.029
47-57	14.467	0.931-224.903	0.056
58 and above			
Gender			
Male	1.049	0.520-2.116	0.893
Female			
Marital Status			
Male	1.988	0.934-4.230	0.075
Female			
Religion			
Christianity	0.772	0.356-1.674	0.512
Islam			
Geopolitical Zone			
Northcentral	1.015	0.282-3.652	0.982
Northeast	1.252	0.350-4.487	0.730
Northwest	4.272	0.568-32.151	0.158
Southeast	3.574	0.917-13.929	0.066
Southwest			
Occupation			
Healthcare workers	0.633	0.199-2.016	0.439
Healthcare support staff			




Fig. 2. How often do you test for latent tuberculosis?

4. DISCUSSION

This survey uses the health belief model (HBM) to determine the factors associated with the willingness to test for latent tuberculosis infection and acceptance of isoniazid by healthcare workers in Nigeria. Our observation revealed that the perception of susceptibility to tuberculosis was low with respect to the probability of getting tuberculosis in the future, family history of tuberculosis, underlying illness, and lifestyle, with the exceptions occupational risk perception (60.9%). Also, the perception of severity of tuberculosis was low. Fortunately, the perceived benefit of tuberculosis vaccination was very high. The finding of this study indicates the need for increase perception of severity and susceptibility to tuberculosis by healthcare workers in Nigeria. This is because as perception to susceptibility and severity increases, the need to accept TB vaccine and other preventive measure such as preventive therapy, regular screening test among population at high risk of occupational exposure to tuberculosis increases [14]. Furthermore, it was noted that high perceived susceptibility and severity to infectious diseases translate into better preventive practice and enhance infections control and prevention [15, 16, 17].

Concerning the perceived severity to tuberculosis, our findings revealed that the fear

of getting tuberculosis differs significantly ($P < 0.05$) among age categories. The group 25-46 years old had the highest proportion that expressed fear of getting tuberculosis. This is maybe because tuberculosis is associated with the middle age group [18, 19] than with the elderly. And it has been shown that age is a predisposing factor to active tuberculosis. regarding the expression of the fear of getting tuberculosis, the result obtained in this study may be mask by the fear of stigmatization. Since in a similar study conducted previously, it was observed that tuberculosis and HIV/AIDS share similar symptoms; and the policy of screening for HIV in tuberculosis patient may generate negative perception [20] and may explain the fear of tuberculosis infections observed among the middle age group in this study. Also, the fear of tuberculosis infections and stigmatization may make people with TB to hide their infection status and the intention to present for screening or diagnosis [21]. In a focus group study [22], in the USA reported that HCWs fears coworker who tested positive for LTBI because they do not know the difference between LTBI and active tuberculosis.

Although, TB incidences and mortality was higher among males than females [1, 23], a higher proportion of the female gender 149 (36.8% vs 26.6%) believed that persons infected with TB could die. This maybe because the

female gender, by her nature (based on African tradition) bears the pain of herself and family members including her husband. By carrying the weight of everybody, she expressed the perception of death due to tuberculosis for her husband and sons.

Our findings showed that most HCWs within the age group 36-46 years has the lowest proportion that agreed that persons infected with TB could die. This is surprising because tuberculosis mortality rate is higher among the age group 35-45 years old [23] compared to the very young and elderly people.

The overall acceptance of LTBI testing was high with nearly four out of every five HCWs expressing the intent to test for LTBI. Although few of the HCWs expressed a definite rejection of latent TBI testing. This proportion is significant given the fact that no one is safe until all are safe. Therefore, there is the need to intensify effort to screen all healthcare workers regularly and educating them on the dangers of tuberculosis. Moreover, the role of healthcare workers as role models in their community is essential in the control and prevention of active tuberculosis and other infectious respiratory diseases [19].

Binary logistic regression analysis found that the demographic characteristics of the respondents were associated with acceptance of LTBI testing. In particular, the findings of this study suggested that the age group 36-46 years old were 14 times more likely to accept LTBI testing. This finding was consistent with a previous study in South Korea in which age was found to be the only risk factor for latent tuberculosis screening among Healthcare Workers [24]. It is recommended that increase tuberculosis education regarding infection control and prevention among the middle age is needed to bring about the desired behavioral change.

Healthcare workers who indicated that they were at high risk of contracting tuberculosis because of their occupation were less likely to undergo screening for LTBI. This finding was surprising because of previous studies in Nigeria and other countries have reported that healthcare workers are at greater risk of tuberculosis infection. Therefore, it is expected that healthcare workers in this study would be more willing to undergo screening for LTBI and accept measures to prevent active tuberculosis.

Concerning who should get testing for LTBI, 115(24.8%) agreed that occupation risk groups and persons in contact with tuberculosis patients should be tested regularly. This finding is consistent with global recommendations for latent tuberculosis testing. The WHO recommended that systematic testing should be performed on people living with HIV(PLWH), adults and child contacts with active tuberculosis cases, prisoners, Healthcare workers, immigrants from high tuberculosis burdened countries, homeless persons, illicit drug users and patients with the following conditions: initiating anti-tumor neurosis factors treatment, receiving dialysis, preparing for transplants, and being affected by silicosis [1]. Thus, identifying and testing people at risk of tuberculosis is an important public health measure to prevent the spread of tuberculosis in the community, hospital or healthcare facility by early diagnosis and initiation of preventive therapy [25].

Interestingly, our finding showed that majority of (56%) of the Healthcare workers in the study never gets tested for LTBI. This is because the policy of systematic or annual LTBI screening has not been implemented for healthcare workers by the National Tuberculosis Control Plan in Nigeria [26].

Regarding isoniazid preventive treatment, majority (87.7%) of the respondents agreed to commence chemoprophylaxis if tested positive for LTBI. Univariable logistics regression analysis revealed that HCWs from the northwest (OR 4.25, 95%CI 1.216-14.688, P=0.023) and southeast (OR 4.414, 95%CI=1.382-14.609, P=0.012) were found to have greater odds of the willingness to accept isoniazid treatment if they are positive for LTBI. The reason for this observation is beyond the scope of this study. It is recommended that further study be carried out to elucidate the factors responsible for the variation in the intention to accept isoniazid in the different geopolitical zones of Nigeria.

Our finding revealed that HBM constructs was associated with the intention to received isoniazid prophylaxis which is in concordance with previous studies [27, 28, 29]. In particular, the finding of this study showed that high perception of seriousness (OR = 2.220, 95% C.I = 1.100-4400 p=0.026) and benefit of tuberculosis immunization (OR = 3.777; 95% C.I = 1.230-11.650; P = 0.020) were the most significant factors that were influencing the intention to accept isoniazid preventive therapy. This finding implies needed for promotional

health messages framing a high risk of tuberculosis and the benefit of carrying out preventive measures.

5. CONCLUSIONS

Most health workers in this study intended to get tested for LTBI and accept isoniazid prevention therapy. The important predictors of the willingness to get tested and accept isoniazid chemotherapy against LTBI include age, geopolitical zone, perception of seriousness and the benefits of tuberculosis immunization. The study has important implication for facilitating government and stakeholders to design policies in Nigeria that will target intervention programmes to enhance tuberculosis prevention and control among healthcare workers and the public. We therefore recommend the inclusion of LBTI testing and treatment in the National TB Control Programme.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declared that no generative AI technology such as large language model (ChatGPT, CAPILLOT ... etc) and text-to image generators have been used during writing or editing of this manuscript.

AVAILABILITY OF DATA AND MATERIALS

All the data supporting this study are included in the article and the supplementary documents.

ETHICS APPROVAL AND CONSENT

Permission to conduct the research was approved by the National Open University of Nigeria (NOUN) research ethic committee. To ensure confidentiality, the names of the participants were not included during data collection, analysis, and presentation. Study participants were given all the information regarding the study and consent to participate was implied by the submission of the completed online survey.

ACKNOWLEDGEMENT

The authors are thankful to the members of the department of Veterinary Public Health and Preventive Medicine and the Department of Veterinary Microbiology, University of Maiduguri and the deputy director, Ms. Lami Daniel of the National Open University of Nigeria for their

efforts and support during the conduct of this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization (WHO), Global tuberculosis report. *Tuberc. Rep.* 2023;13–14.
2. Dutta NK, Karakousis PC. Latent tuberculosis infection: myths, models, and molecular mechanisms. *Microbiology and Molecular Biology Reviews.* 2014 Sep;78(3):343-71.
3. Berry MP, Graham CM, McNab FW, Xu Z, Bloch SA, Oni T, Wilkinson KA, Banchereau R, Skinner J, Wilkinson RJ, Quinn C. An interferon-inducible neutrophil-driven blood transcriptional signature in human tuberculosis. *Nature.* 2010 Aug 19;466(7309):973-7.
4. Day CL, Abrahams DA, Harris LD, van Rooyen M, Stone L, de Kock M, Hanekom WA, Hiv-1 infection is associated with depletion and functional impairment of mycobacterium tuberculosis-specific cd4 t cells in individuals with latent tuberculosis infection. *J. Immunol.* 2017;199:2069–2080.
5. El-Mokhtar MA, Elgendy SG, Eldin AS, Hassan EA, Hasan AAA, Abdel Hameed MR, Sayed D, Salama EH. Hepatitis C virus affects tuberculosis-specific T cells in HIV-negative patients. *Viruses.* 2020 Jan 15;12(1):101. DOI: 10.3390/v12010101. PMID: 31952232; PMCID: PMC7019953
6. Mohammed D, Khallaf S, El-Naggar M. et al. Interleukin-10: A potential prognostic marker in patients with newly diagnosed multiple myeloma. *Research in Oncology.* 2021;17(1):38-41. DOI: 10.21608/resoncol.2021.51503.1127
7. Geldmacher C, Schuetz A, Ngwenyama N, Casazza JP, Sanga E, Saathoff E, Boehme C, Geis S, Maboko L, Singh M et al. Early depletion of mycobacterium tuberculosis-specific t helper 1 cell responses after hiv-1 infection. *J. Infect. Dis.* 2008;198:1590–1598.
8. World Health Organization (WHO), Global tuberculosis report, WHO Library Cataloguing-in-Publication Data; 2016.

9. Houben RM, Dodd PJ. The global burden of latent tuberculosis infection: a re-estimation using mathematical modelling. *PLoS medicine*. 2016 Oct 25;13(10): e1002152.
10. Apriani L, McAllister S, Sharples K, Alisjahbana B, Ruslami R, Hill PC, Menzies D. Latent tuberculosis infection in healthcare workers in low-and middle-income countries: an updated systematic review. *European Respiratory Journal*. 2019 Apr 1;53(4).
11. Umo AN, Asuquo AE, Abia-Bassey LN, Moses AE. Prevalence of latent tuberculosis infection among health workers resident in Akwa Ibom State, South-South Nigeria. *IJTDH*. 2016;12(3):1-7.
12. Ejeh FE, Saidu AS, Owoicho S, Maurice NA, Jauro S, Madukaji L, Okon KO. Knowledge, attitude, and practice among healthcare workers towards COVID-19 outbreak in Nigeria. *Heliyon*. 2020 Nov 1;6(11).
13. ECOM-EM. Standard questionnaire on risk perception of an infectious disease outbreak This tool contains example questions for public surveys on risk perception of (an outbreak of) an infectious disease; 2015.
14. Wong LP, Alias H, Wong PF, Lee HY, AbuBakar S. The use of the health belief model to assess predictors of intent to receive the COVID-19 vaccine and willingness to pay. *Human vaccines & immunotherapeutics*. 2020 Sep 1;16(9):2204-14.
15. Alqudeimat Y, Alenezi D, AlHajri B, Alfouzan H, Almokhaizeem Z, Altamimi S, Almansouri W, Alzalalah S, Ziyab AH. Acceptance of a COVID-19 vaccine and its related determinants among the general adult population in Kuwait. *Medical Principles and Practice*. 2021 Jan 22;30(3):262-71.
16. Gan L, Chen Y, Hu P, Wu D, Zhu Y, Tan J, Li Y, Zhang D. Willingness to receive SARS-CoV-2 vaccination and associated factors among Chinese adults: A cross sectional survey. *International Journal of Environmental Research and Public Health*. 2021 Feb;18(4):1993.
17. Lin C, Tu P, Terry TC. Moving the needle on racial disparity: COVID-19 vaccine trust and hesitancy. *Vaccine*. 2022 Jan 3;40(1):5-8.
18. Narasimhan P, Wood J, MacIntyre CR, Mathai D. Risk factors for tuberculosis. *Pulmonary medicine*. 2013;2013(1): 828939.
19. Ejeh FE, Undiandeye A, Okon K, Kazeem HM, Kudi AC. Isolation and immunological detection of Mycobacterium tuberculosis from HIV and non-HIV patients in benue state, Nigeria. *Ethiopian Journal of Health Sciences*. 2020 Jan 1;30(1).
20. Ntinginya NE, Te Brake L, Sabi I, Chamba N, Kilonzo K, Laizer S, Andia-Biraro I, Kibirige D, Kyazze AP, Ninsiima S, Critchley JA. Rifapentine and isoniazid for prevention of tuberculosis in people with diabetes (PROTID): Protocol for a randomised controlled trial. *Trials*. 2022 Jun 10;23(1):480.
21. Nyasulu P, Sikwese S, Chirwa T, Makanjee C, Mmanga M, Babalola JO, Mpunga J, Banda HT, Muula AS, Munthali AC. Knowledge, beliefs, and perceptions of tuberculosis among community members in Ntcheu district, Malawi. *Journal of Multidisciplinary Healthcare*. 2018 Aug 16:375-89.
22. Joseph HA, Shrestha-Kuwahara R, Lowry D, Lambert LA, Panlilio AL, Raucher BG, Holcombe JM, Poujade J, Rasmussen DM, Wilce M. Factors influencing health care workers' adherence to work site tuberculosis screening and treatment policies. *American Journal Infectious Control* 2004;32:456–461.
23. Gichuki J, Mategula D. Characterisation of tuberculosis mortality in informal settlements in Nairobi, Kenya: Analysis of data between 2002 and 2016. *BMC Infectious Diseases*. 2021 Dec; 21:1-8.
24. Jong Lee K, Ae Kang Y, Mi Kim Y, Cho SN, Wook Moon J, Suk Park M, Kyu Kim S, Chang J, Sam Kim Y. Screening for latent tuberculosis infection in South Korean healthcare workers using a tuberculin skin test and whole blood interferon- γ assay. *Scandinavian journal of infectious diseases*. 2010 Sep 1;42(9):672-8.
25. Ayub A, Yale SH, Reed KD, Nasser RM, Gilbert SR. Testing for Latent Tuberculosis. *Clinical Medical Research*. 2004;2:191.
26. Federal Ministry of Health,. The national strategic plan for tuberculosis and leprosy control 2012. [WWW Document]. NTBLCP. URL; 2010-2015. Available:www.healthsystems2020.org

27. Cavanaugh JS, Modi S, Musau S, McCarthy K, Alexander H, Burmen B, Heilig CM, Shiraishi RW, Cain K. Comparative yield of different diagnostic tests for tuberculosis among people living with HIV in Western Kenya. PloS one. 2016 Mar 29;11(3): e0152364.
28. Lee JY, Park JY, Kim MS, Kim JH, Lee JY. Selection of health care workers for screening for latent tuberculosis infection. Journal of Tuberculosis Research. 2019 Jun 19;7(02):65.
29. Thomas TA. Tuberculosis in children. Thoracic surgery clinics. 2019 Feb 1;29(1):109-21.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/122966>