



Transposition of Abdominal Inferior Vena Cava to the Left: A Rare Case Report

M. V. Ravishankar^a, N. B. Pushpa^a,
Prasanna K Santhekadur^b and Bhagath Kumar Potu^{c*}

^a Department of Anatomy, JSS Medical College, JSS Academy of Higher Education and Research, Mysuru, Karnataka, India.

^b Department of Biochemistry, JSS Medical College, JSS Academy of Higher Education and Research, Mysuru, Karnataka, India.

^c Department of Anatomy, College of Medicine and Medical Sciences, Arabian Gulf University, Manama, Bahrain.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2023/v21i4807

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96572>

Case Report

Received: 18/12/2022

Accepted: 24/02/2023

Published: 08/03/2023

ABSTRACT

Development of the vascular system in humans is a complex process that involves formation, retention, regression, and reconnection of primitive vascular channels, ultimately forming a definite pattern of the venous vascular architecture. Any errors in these processes would lead to anomalous venous structures, which would pose severe diagnostic and therapeutic challenges. Herein, we report a case of left-sided inferior vena cava that was noticed in the posterior abdominal wall of a male cadaver aged about 80 years. Although venous anomalies are common in their occurrence, a left-sided transposition of the inferior vena cava is a rare and significant finding in

*Corresponding author: E-mail: potubk@agu.edu.bh;

retroperitoneal surgeries because this can be misdiagnosed as para-aortic lymphadenopathy, a tumor or dilated gonadal vein that may result in iatrogenic damage during surgeries. The unusual location and course of the inferior vena cava and its embryological basis are discussed in this case.

Keywords: *Left-sided inferior vena cava; venous transposition; venous anomalies; inferior vena cava clinical implications.*

1. INTRODUCTION

The inferior vena cava (IVC) is a large vein normally located on the right side of the vertebral column beside the aorta on its left. It is formed at the level of L₄ vertebrae by the union of the right and left common iliac veins [1]. The formation of IVC is a complex process that involves the vitelline vein, supracardinal, subcardinal, and posterior cardinal veins for its definitive and uniform morphological existence. The occurrence of a left-sided IVC is the second most common anomaly after duplicated IVC [2]. These infrequent variations of IVC, which are incidentally found during diagnostic interventions which may not stand solitary but may tend to show some associated variation in other vascular channels, such as ovarian or testicular (gonadal), suprarenal, and renal venous channels, respectively [3]. A thorough assessment of structural anomalies, such as transposition of the IVC, needs to be understood from prophylactic, diagnostic, and therapeutic standpoints. This

case report involved a left-sided transposition of the IVC found during routine anatomy dissection sessions.

2. CASE DESCRIPTION

During the routine dissection of the posterior abdominal wall contents in a male cadaver aged about 80 years, we noticed an IVC located on the left side of the abdominal aorta (AA) possessing all the tributaries in its regular anatomical pattern (Fig 1). Upon further examination, it was found that the part of IVC was showing a disposition to the left just below the origin of the superior mesenteric artery. It later continued ascending further on crossing the AA as an infra hepatic part of its course followed by a retrohepatic location. The left-sided IVC finally entered the thorax at the level of the 8th thoracic vertebrae through the vena cava opening of the diaphragm. No other gross anomalies were noted in this case.




Fig. 1. Showing left-sided transposition of the inferior vena cava: 1: left-sided inferior vena cava; 2: abdominal aorta; 3: Right kidney; 4: Left kidney; 5: Right renal vein; 6: Left renal vein; 7: Superior mesenteric artery

3. DISCUSSION

Normal development of the IVC occurs between the sixth to tenth weeks of gestation [4]. This is the period during which all the anomalies of IVC are expected to take place. The embryogenesis of the human circulatory system is regulated by many molecular signaling pathways and genes. Studies reported that the mutations in these genes, particularly *TIE3* gene mutations might cause venous channel malformations [5]. Embryologically, IVC is formed by the right supracardinal vein. In case of left-sided IVC as seen in the present case, the left supracardinal vein persists and the right supracardinal vein regresses. In cases of left supracardinal vein regression failure, retention of another parallel segment of the venous channel gives the appearance of IVC duplication in the posterior abdominal wall. In our case, a left-sided IVC was found lying beside the AA without any duplicating remanent channel on the right side. Hence no duplication pattern was found.

Caval anomalies are sporadic, often they are incidental findings of radiological investigations. A rare case of left-sided IVC was encountered during the post-mortem study of a 36-year-old woman, who died due to accidental trauma. The IVC was showing its unusual origin from the point of confluence of the right and left iliac veins lying on the left side of the AA. IVC possessed its regular tributaries like right and left renal veins. The IVC also showed its contralateral course exactly at the point of drainage of its left renal tributary, and its further course was lying in the infra hepatic or supra renal region of the posterior abdominal wall. These findings are consistent with our present study findings encountered during cadaveric dissection [2]. A systematic review conducted on left-sided IVC from 73 different studies revealed that the prevalence of IVC is as low as 0.1-0.4% showing a slight male preponderance [6]. Though these anomalies of IVC are rare and asymptomatic, deep vein thrombosis (DVT) is one of the most commonly associated risk factors which can be clinically correlated with the incidence of transposition of IVC. Its displacement and altered course may increase the susceptibility for thromboembolism due to undue compression of its venous channels [7]. It is also very important for surgeons to be aware about left-sided IVC during complicated retroperitoneal surgical procedures such as abdominal aortic aneurysm repair, left-sided nephrectomy, oblique lumbar fusion or IVC filter placement, and

sigmoidectomy [8-11]. However, these surgical complications of left-sided IVC can be avoided with an accurate pre-operative evaluation of the retroperitoneal venous system under radiological interventions such as computed tomography (CT) or magnetic resonance imaging [12] (MRI).

4. CONCLUSION

Major deep venous channel anomalies are relatively less common when compared with superficial veins. It is very important to identify eccentric anatomical variations of large veins from the perspective of clinical, diagnostic, and therapeutic standpoints to avoid fatal consequences in medical practice.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

PERMISSION

Obtained for cadaver dissection, presentation, and research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Standing S. Gray's anatomy: The anatomical basis of clinical practice. 41st ed. New York; Elsevier; 2016.
2. Rajabnejad Y, Aliakbarian M, Rajabnejad A, Motie MR. Left sided inferior vena cava encountered during organ retrieval surgery: Report of two cases. *Int J Organ Transplant Med.* 2016;7:229-232.
3. Ghandour A, Partovi S, Karuppasamy K, Rajiah P. Congenital anomalies of the IVC—embryological perspectives and clinical relevance. *Cardiovasc Diagn Ther.* 2016;6:482-492.
4. Giordano JM, Trout III HH. Anomalies of the inferior vena cava. *J Vasc Surg.* 1986;3:924-928.
5. Kim JS, et al. The molecular pathophysiology of vascular anomalies: Genomic research. *Arch Plast Surg.* 2020;47:203-208.

6. Ang WC, Doyle T, Stringer MD. Left-sided and duplicate inferior vena cava: a case series and review. Clin Anat. 2013;26:990-1001.
7. Zhang L, Guan WK. Deep vein thrombosis in patient with left-sided inferior vena cava draining into the hemiazygos vein: A case report. World J Clin Cases. 2021;9:672-676.
8. Wu WK, Guth CM, Mulherin JL. Abdominal aortic aneurysm repair requiring left-sided inferior vena cava interposition. Ann Vasc Surg. 2020;68:573–575.
9. Wang LH, et al. Left transperitoneal laparoscopic partial nephrectomy in the presence of a left-sided inferior vena cava. Urology. 2011;78:469–473.
10. Berry CA. Oblique lumbar interbody fusion in patient with persistent left-sided inferior vena cava: Case report and review of literature. World Neurosurg. 2019;132:58–62.
11. Matsunaga M, Fukahori M, Ushijima T, Miwa K. Left-sided inferior vena cava. Case Reports. 2015:bcr2015211268.
12. Li SJ, Lee J, Hall J, Sutherland TR. The inferior vena cava: anatomical variants and acquired pathologies. Insights Imaging. 2021;12:123.

© 2023 Ravishankar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96572>