



## **Advanced Physiotherapy Rehabilitation in Patient with Low Back Pain- A Case Report**

**Alpriya Pathan <sup>a≡</sup>, Ragini Dadgal <sup>a∞\*</sup>, Neha Chitale <sup>a#</sup>, Rakesh Kovala <sup>a†</sup>  
and Irshad Qureshi <sup>a‡</sup>**

<sup>a</sup> Department of Physiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha, Maharashtra, 442001, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i60B34921

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77827>

**Case Study**

**Received 10 October 2021**  
**Accepted 13 December 2021**  
**Published 26 December 2021**

## **ABSTRACT**

Low back pain is a common condition due to many possible causes. The causes include improper lifting, improper posture, lack of regular exercise and fracture. Low back pain is experienced by everyone at least once in their lifespan. Proper management of low back pain is very important to prevent further complications. Low back pain can be managed by medication, surgery or by physiotherapy depending upon the cause. Physical therapy is one of the treatment for various low back pain. To manage low back pain the cause plays a major role, the therapist should take some efforts to know the cause and treat accordingly. Manual therapy is very useful for improving range of motion and tightness of the muscle.

**Keywords:** Low back pain; rehabilitation; physiotherapy.

### **1. INTRODUCTION**

Low back pain is a common condition due to many possible causes. The causes include

improper lifting, improper posture, lack of regular exercise and fracture [1]. Low back pain is experienced by everyone atleast once in there lifespan. Proper management of low back pain is

<sup>≡</sup>BPT Student;

<sup>∞</sup>Assistant Professor;

<sup>#</sup>MPT Student;

<sup>†</sup>Associate Professor;

<sup>‡</sup>Professor;

\*Corresponding author: E-mail: dr.raginidadgal@gmail.com;

very important to prevent further complications. Low back pain can be managed by medication, surgery or by physiotherapy. Depending upon the cause physiotherapy approach differs.

Postural and traumatic back pain is common. Low back pain may be acute or chronic. Acute low back pain is the pain which lasts for few days to few months, chronic low back pain is the pain which lasts for more than 6 months [2]. Acute pain related to definite episode of trauma or spontaneous in onset and chronic low back pain it is due to inflammation or mechanical alterations like inclined posture [3]. Hyperlordosis and lordosis are postural effect on the spine [4]. Low back pain recurrence percentage of one year is 54 percent to 90 percent [2]. The goal of physical therapist is to assess and rehabilitate the patient. In a study performed by Rekha Chaturvedi, et al; showed significant improvement in pain in group that received core strengthening exercises. Therefore, core strengthening exercises might be recommended for reducing pain in patients with chronic low back pain. Which helped us to set long term goals.

## 2. PATIENT INFORMATION

The patient was 47 years old female who had been experiencing low back pain from 1 year and she is farm worker. The patient was very limited in activities of daily living and she was unable to walk for long time as she had pain in both legs which was radiating in nature. Tingling sensation on both legs occur due to farm work. The patient reported no significant orthopedic issue. Tingling and numbness in tarsal and metatarsal region is present.

## 3. CLINICAL PRESENTATION

She was examined in supine with both the shoulders in level. Upon inspection, the patient maintains both legs abducted and slightly internally rotated, knee extended with pillow support below both knees, legs and ankle, both ankles slightly plantar flexed. She complained of dull aching pain which was 7/10 on numerical pain rating scale (NPRS) on rest and 9/10 on standing or unsupported sitting for 10 minutes with Oswestry Disability Index (ODI) (with

permission) score of 80% disability. On observation there is no swelling, no deformity. Range of motion and manual muscle testing was done.

### 3.1 Activities of Daily Living

Patient specific functional scale was used to measure functional outcome. 5 activities was selected and the score was 4/10.

### 3.2 Therapeutic Intervention

Treatment starts on the first day of the visit to the hospital after filling consent form from the patient. The primary goal was to reduce the pain of the patient and to train the patient for activities of daily living.

**Day 1-14:** Hydrocollateral pack was given for 10 minutes for instance heat increases joint temperature, improves the circulation and helps in the muscle relaxation and muscle spasm and helps in relaxing the tightness of the muscle. Interferential therapy was set up in the treatment protocol, If it helps in pain relief and increases the local blood flow of that area and relieves the oedema also. Application of Interferential therapy is for only for 10 minutes in four pole method and intensity depends on patient tolerance level. Traction is also given to the patient for low back pain to treat the condition of the spine.

**Day 14-30:** The aim was to maintain the goals achieved in first 14 days and to strengthen the back and lower limb muscles for which exercise were given.

Exercises like back extension exercises, straight leg raise, pelvic bridging. Each exercises should perform for 2 sets of 10 repetitions.

Back extension exercises can strengthen the lower back muscles which include erector spinae. Straight leg exercise helps in assessment of sciatic pain and for herniated disc. Pelvic bridging exercises help in trunk stability and increases muscular strength. This protocol should follow for a week. Avoid bending and heavy lifting activities.

**Table 1. Range of motion**

Joint	Right	Left
Hip flexion	0 to 120	0-110
Hip extension	25	20

**Table 2. Manual muscle testing**

<b>Joint</b>	<b>Right</b>	<b>Left</b>
Hip flexion	Grade 3	Grade 3
Hip extension	Grade 4	Grade 4
Trunk forward flexion	Grade 3	
Trunk extension	Grade 4	

**Day 30-60:** The patient was continued with the exercised given in initial phase and core strengthening was started.

**Core Strengthening:** For static exercises (Each exercise was repeated for 10 times with hold time of 10 sec in one set.)

The subjects were asked to hold the core muscle by drawing in maneuverer.

The subjects were asked to perform abdominal drawing-in manoeuvre [5].

**Quadrupedal stance (Bird dog exercise):** Subjects were required to remain in a quadruped position with both hands and knees flat on the surface; a leg and the contralateral arm were raised at a moderate velocity in horizontal position. **Cross curl-ups:** Subjects were asked to lie in a supine position, hands folded behind the back, elbows pointed towards the sides, knees in a flexed position, feet resting on a mat; subjects rolled up until the scapulae left the mat, then rotated at a mild velocity to the left and right.

### 3.3 Outcomes Measures

**Numeric Pain Rating Scale:- Pre 7/10, Post 3/10.**

**Modified Oswestry Scale: - Pre 60% disability, Post – 20% disability.**

**Table 3. Range of motion post 2 months of intervention**

Joint	Right	Left
Hip flexion	0 to 160	0-150
Hip extension	35	30

**Table 4. Manual muscle testing post 2 months of intervention**

Joint	Right	Left
Hip flexion	Grade 4	Grade 4
Hip extension	Grade 4	Grade 4
Trunk forward flexion	Grade 4	
Trunk extension	Grade 5	

### 4. DISCUSSION

The patient was 47 years old female who had been experiencing lbp from 1 year ago and she is farm worker. The patient was very limited in activities of daily living and she was unable to walk for long time as she had pain on both legs .radiating pain occur on both legs tingling sensation on both legs occur due to farm work she has to go to farm daily but cannot able to walk for long time.

Specific exercise training is necessary for chronic low back pain to produce functional improvement [5]. Various manual therapy should be given. Hydrocollateral pack is used for superficial heating which relaxes the muscles and improves blood circulation while interferential therapy is a technique which combines two low frequency current to make a medium frequency current

which also helps in reducing the pain [6]. There is impact of low back pain on functional activities. It affect on activities a lot. And while treating the patient fear and pain are the two main factors and kinesiophobia is one of the factor in the patient. The patient get scared of the doing movements. So Tampa kinesiophobia scale are used for acute and chronic low back pain. Kinesiophobia creates a very adverse effects on the patient. Things should keep in mind that patient should not get panic.and and while giving exerces the patient should not have anxiety while doing exercises [7] and kinesiophobia it act on acute and chronic low back pain it effect on both [8].

There are so many surveys to check the kinesiophobia. And its rating will measure the outcomes plays a very crucial role in the low back pain. kinesiophobia is more in people who

don't do exercises because of this it create phobia in people and the people who exercises they have less experience of kinesiophobia [7]. Sensory testing and reflexes are the components of physical examination draws attention to radiculopathy pain [9].while applying stretching technique proper duration and frequency are required.

A systemic review on non- specific low back pain done by Rebecca Gordon; et al; found that improving the flexibility of the muscle-tendon and ligaments in the back increases the range of motion and assists with the patient's functional movement. Increasing core muscular strength can assist in supporting the lumbar spine. Which was mainly achieved by ART as it simultaneously locating and breaking down of scarred tissue and adhesion, since that is the primary cause of pain, stiffness, weakness, as well as physical dysfunction that are typically associated with soft tissue injuries.

## 5. CONCLUSION

Low back pain is very common condition in mostly in females. The clinical course of low back pain presented in general practice, for the most patients, clearly is less favorable than favorable than looking forward too, it takes more than just a few weeks to relapses and recover takes place within a year in most cases. Manual therapy are very useful for improving range of motion and tightness of the muscle. Physiotherapy plays an important role in treating low back pain. Regular physiotherapy treatment is needed.

## CONSENT AND ETHICAL APPROVAL

As per international standard or university standard guideline Patient's consent and ethical approval has been collected and preserved by the authors.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Holy D, Brooks P, Blyth F, Buchbinder R, the epidemiology of low back pain Best Practices Clin Rheumatol Rheumatol. 2010;24(6):769-81.
2. Frost H, Lamb SE, Doll HA, Carver PT, Stewart-Brown S. Randomised controlled trial of physiotherapy compared with advice for low back pain. BMJ. 2004; 329(7468):708.
3. Moseley L. Combined physiotherapy and education is efficacious for chronic low back pain. Aust J Physiother. 2002;48(4): 297–302.
4. Brumagne S, Janssens L, Janssens E, Goddyn L. Altered postural control in anticipation of postural instability in persons with recurrent low back pain. Gait Posture. 2008;28(4):657-62.
5. Deshmukh MK, Phansopkar PA, Kumar K. Effect of muscle energy technique on piriformis tightness in chronic low back pain with radiation. J Evol Med Dent Sci. 2020;9(44):3284–9.
6. Naqvi WM, Vaidya L, Kumar K. Impact of low back pain on fear of movement and functional activities. International Journal of Research in Pharmaceutical Sciences. 2020;11:4830-35.
7. Goyal CV, Naqvi WM. Lordoscoliosis and hyperlordosis in quadriplegic cerebral palsy. The Pan African Medical Journal. 2020;36.
8. Bais AS, Mishra SA, Darda PP, Naqvi WM. Physiotherapy rehabilitation in sensory motor polyneuropathy along with lumbar radiculopathy: A novel case report. European Journal of Molecular & Clinical Medicine. 2020;7(11):2653-68.
9. Bais A, Bawiskar D, Naqvi WM, Sahu A. A case study on the impact of physiotherapy on unilateral foot drop after lumbar fusion and discectomy. Medical Science. 2020; 24(103):1773-9.

© 2021 Pathan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/77827>