



Role of Bamboo Forest for Mitigation and Adaptation to Climate Change Challenges in China

Regassa Terefe^{1,2,3*}, Liu Jian^{1,2} and Yu Kunyong^{1,2}

¹*University Key Laboratory for Geomatics Technology and Optimize Resources Utilization in Fujian Province, Fuzhou, Fujian 350002, China.*

²*College of Forestry, Fujian Agriculture and Forestry University, Fuzhou, Fujian 350002, China.*

³*Oromia Agriculture Research Institutes, Bako Agriculture Research Center, P.O.Box 03, West Shoa, Oromia, Ethiopia.*

Authors' contributions

This work was carried out in collaboration among all authors. Among RT designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors LJ and YK managed the literature part of the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JSRR/2019/v24i130145

Editor(s):

(1) Dr. Rajvir Yadav, Professor, Department of Farm Engineering, Junagadh Agricultural University, Junagadh, India.

Reviewers:

(1) Dickson Adom, Kwame Nkrumah University of Science and Technology, Ghana.

(2) Antonio L. Beraldo, Campinas University, Brazil.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/44239>

Review Article

Received 20 July 2018
Accepted 27 September 2018
Published 20 July 2019

ABSTRACT

Bamboo is one of the fastest growing plants on the planet, with many attributes which make it a useful potential resource for humankind. Though having fast growth and good regeneration performance after harvesting is a unique characteristic of the specie. It enhances a high carbon storage potential particularly when the harvested culms are transformed into durable products. China has many bamboo species with distribution and area coverage's, and highly connected in using the production of bamboo resources. Its characteristics make it an ideal solution for the environmental and social consequences of tropical deforestation. This review paper aims to assess the contribution of bamboo in mitigating and adapting impacts of climate change and its importance regarding ecological and socio-economic benefits. The review summarised the role of bamboo forests towards mitigating and adapting its potential to overcome the impacts of climate change currently seen globally and particularly to China. Therefore, advancing bamboo farming systems at different levels, it's advantages to reduce greenhouse gas in the atmosphere and expanding bamboo forests in future under wider use and intensive management is recommended.

*Corresponding author: E-mail: regassaterefe@yahoo.com;

Keywords: Bamboo; carbon sink; climate change; greenhouse gas.

1. INTRODUCTION

Bamboo is a grass type of Gramineae family, and it is an important component of many forest environments. It adapts easily to a range of climatic and soil conditions, and is widely distributed in the tropical and subtropical zones approximately between 46° N and 47° S latitude, covering a total area of about 31.5 million ha, and accounted for about 0.8% of the World's total forested area in 2010 [1]. It has unique features that distinguish it from most other woody plants. For example culms that are connected by an extensive system of rhizomes, leading to emerging new culms by rapid asexual reproduction [2,3,4].

The bamboo resources are distributed in many countries in the World, majorly found in Asia, Africa, and Latin America; however, the origin of most of the species lies in Southeast Asia. Worldwide bamboo families categorised with more than 107 genera and 1300 species [5,6]. Along with this, all China has the highest bamboo species diversity, with 39 genera and 509 species, accounting for 36% and 39%, respectively, of the total bamboo genera and species with the rest of the World [6]. Besides that, China's bamboo forests cover an area of 4.84 million ha in 2005, which accounts for 2.8% of China's forested area and 15.4% of the World's area of bamboo [7]. Bamboo forests are the most important non-wood forest product and substitute wood products. As a resource, bamboo forests are an important part of ecosystems, which provide some basic environmental services. Bamboo provides food and raw materials, reduces water erosion on slopes, regulates water flows, and it acts like windbreak in shelterbelts which offer protection against storms [1]. In addition to this because of its special root re-sprouting regeneration strategy bamboo forests, it generates a good potential for carbon storage mechanisms, and water and soil conservation and much more advantageous [5].

Many studies have shown that well managed and regularly harvested bamboo can sequester more carbon than bamboo in a natural state. Despite that, it can sequester more carbon than fast-growing tropical and subtropical trees under comparable conditions. If bamboo forests were not managed through annual harvesting practices, they would be significantly less

effective in carbon sequestration. This review paper aims to assess the contribution of bamboo in mitigating and adapting impacts of climate change and its importance regarding ecological and socio-economic benefits.

2. ROLES OF FORESTS IN MITIGATING CLIMATE CHANGE

Forests are one of the biggest reservoirs of carbon, so they help to keep the carbon cycle and other natural processes working and help reduce climate change. However, forests can also be one of the most significant sources of CO₂ emissions [8]. Also, forests provide a wide range of ecological, social, and economic benefits, in the form of goods and services to society, that is much less easy to quantify. Besides that, the demand for timber and related products will require more efficient and sustainable use of natural resources. Forests are the most vulnerable climate dependent systems but have also been recognised to have a significant and crucial contribution to addressing the challenges of mitigation and adaptation in tandem with the issues of livelihoods, economic growth and development. However, the most recent report from the International Union of Forest Research Organizations (IUFRO) indicates the gloomy picture about the future of the World forests in a changing climate. While, it suggests that in a warmer world, the current carbon regulating services of forests as carbon sinks may be entirely lost as land ecosystems could turn into a net source of carbon dioxide later in the century [9]. It plays in combating climate change impact through reducing the emissions from deforestation and forest degradation has become a fundamental issue to international dialogues on preventing the current global temperature increases [10]. The challenges of climate change seen in affecting forest ecosystems in their structure and morphology, thus causing an implication on the functionality of forests in every corner of the World [11, 12]. Besides that, it is considered one of the greatest threats facing humanity in the current global situations. According to the Intergovernmental Panel on Climate Change (IPCC), global warming is unambiguous, with evidence towards the increases of average air and ocean temperatures, which leads to aggravating melting of snow and ice and sea levels [8,13].

3. BAMBOOS FOR CLIMATE CHANGE ADAPTATION

Bamboos are one of the World's strongest and fastest growing woody plants capable of providing ecological, economic and livelihood security to the people, distributed over ranges of climate from mild-temperate to tropical. Bamboo's fast growth ability to grow on varied soils and climate, renewability and positive socio-economic impacts make them an excellent alternative for combating climate change [14]. The high growth potential and ability to store large amounts of carbon make sequestration, and on the other hand, their environmental and socio-economic services can help communities in developing countries to adapt to climate change impacts. According to the research result by International Network of Bamboo and Rattan (INBAR) had shown that well-managed bamboos could be an effective in carbon sink and perform better than other species like Chinese fir and eucalyptus growing under similar conditions. Furthermore, the necessities of bamboo management with a period, sustainable way and selective harvesting mechanism of the stem which are changed into products that can hold carbon for many years. The increasing popularity of durable bamboo products ensures that for the foreseeable future, productive bamboo systems can be considered as an important carbon sink [15].

3.1 Bamboo for Timber Demand and Climate Change

The demand for timber and different agricultural products will continue to increase with the global population. Instead of satisfying the increasing demand for different commodities, the global policies must need to shift towards using efficient and sustainable production systems [13]. Bamboo is one of an alternative resource that used to play an important contribution towards reducing direct pressure on forest resources [13, 24]. Furthermore, one of the best example in China, following the logging bans of certain forests resources came into effect in 1998, while bamboo forests have been used as a possible substitute for timber.

Magel et al. [16] argues that the growth of new shoots in bamboo plantations occur because of the transfer of the energy accumulated in culms through photosynthesis in the previous year. The result of such growth of bamboo culms is not driven by its own carbon sequestration, but by sequestration in previous seasons in other parts

of the bamboo system, and such growth of new shoots is not an indicator of sequestration rate. Another report by Zhou [17] show that as the bamboo system requires more inputs in the shooting period of young culms, which means when new shoots developed during that time the high growth of bamboo shoots, can be equated with a high rate of carbon sequestration. The maturity period of most bamboo culms estimated between 7-10 years, approximately, after that they deteriorate rapidly, releasing carbon from the aboveground biomass back into the atmosphere [18]. Therefore, at a natural circumstance, bamboo will reach a stable level of above ground carbon relatively quickly, even though carbon accumulation through sequestration is offset by carbon release through deterioration of old culms.

3.2 Suitable Ecological Growing Conditions for Bamboo Species

Naturally, bamboo species has the potential to grow at different altitudinal range from 0 to 4000 meter above sea level. It prefers well-drained sandy loams to loamy clay types of soils originated from river alluvium or underlying rock. In most of the bamboo thrive well at annual average temperatures range of 8.8 – 36°C and an annual rainfall of 1270 – 4050 mm. On the other hand, some bamboo species are also growing under high rainfall areas, while some can tolerate limited winter frost [19]. ICFRE [20] reported as one of the fastest growing species in the planet, under the ideal environment; it can be growing up to one meter a day. The biomass of freshly planted bamboo plantation increases rapidly for the early six to eight years after which emergence and death of culms tend to become equal.

4. MANAGEMENT AND HARVESTING OF BAMBOO

Bamboos have well-developed rhizomes with good root systems which help to obtain strengthen during their existence. Its culms mature within three to four years and naturally die after eight to ten years, if not used the products [21]. The periodic removal of mature culms from each bamboo clump and this cycle of removal may vary from two to four year. Thus, provide a highly renewable resource with a high degree of sustainability. It makes bamboos acquiescent to sequester accumulated CO₂ from the atmosphere throughout the lifetime [22]. The sequestration rate of bamboo is higher during the initial eight to ten years period of fast growth.

Numerous studies categorize the production management of bamboo management practices in different five major points: timber, shoot, pulp, ornamental and water/soil conservation benefits.

5. BAMBOOS FOR CLIMATE CHANGE MITIGATION

Bamboo grows more rapidly than any other trees and reaches to give yield within three to four years after planting. Hence, it is one of the fast growing and responding well against, a drought which can make the species more acceptable in making the evergreen environment in addition to soil and water conservation, carbon storage and rehabilitation of degraded lands [23]. It offers one of the quickest ways to remove vast amounts of that CO₂ from the atmosphere. Lou et al. [24] report that at the age of 9 -10 years old in moso bamboo (*Phyllostachys pubescens*) plantation the above ground carbon stock ranges between 25 to 32t ha⁻¹ in China. Furthermore, another study shows that in *P. pubescens* and *P. bambusoides* from natural forests in Japan have an aboveground carbon stock of 78.6 t ha⁻¹ and 52.3 t ha⁻¹, respectively [25]. A four-year mixed bamboo plantation (*Bambusa vulgaris*, *B. balcooa*, and *B. cacharensis*) in India shows that the aboveground carbon stock is about 61.05t ha⁻¹ [26]. Despite that, it provides a minimum CO₂ gas and generates up to 25% more oxygen than other trees within the same level. One hectare of bamboo can sequester up to 62 t of CO₂ yr⁻¹, whereas equivalent of young forest sequesters 15 t of CO₂ yr⁻¹. The *Guadua* plantations in Costa Rica estimated to absorb 17 t of CO₂ ha yr⁻¹ as the study showed by Janssen [27]. Another research study by INBAR states that over the past 15 years, areas under bamboos in Asia grew by 10%. Studies have estimated that the carbon stored in Chinese bamboo forests will increase from 727.08 Tg C in 2010 to 1,017.54 Tg C in 2050, which equates to an increase of nearly 40% in 40 years. This represents a significant contribution to the Chinese forest carbon stock and a range that shows that policies aiming at combating climate change with bamboos can indeed have significant promise [28]. For example: by INBAR's, as modelling shows that a managed moso bamboo forest accumulates about 300 t of carbon ha⁻¹ after 60 years. As well, it does also produce more biomass under well managed and regular harvesting of mature culms. Another report by Lou et al. [24], confirmed that the amount of carbon sequestration between a fast-growing Chinese Fir plantation and monopodial

(*P. pubescens*) plantation modelled for subtropical agro-ecologies in South East China and the results showed that, bamboo sequestered more carbon than the Chinese Fir in the first 5 years. This might be due to rapid early growth; bamboos sequester more carbon in the early years of a plantation than comparable forest trees. In the other way, unmanaged bamboo stands do not store high levels of carbon, as their productivity is low and the accumulated carbon returns quickly to the atmosphere as the older culms decompose [28].

5.1 Carbon Sequestration Potential of Bamboo

Carbon sequestration is the process of capturing and storing atmospheric carbon dioxide. Naturally, the performance of bamboo plantation is different comparing with other tree species since it has a fast growth rate with high re-growth behaviour after harvesting. Despite that, it has a high carbon storage potential according to Zhou and Jiang studies [29], especially when the harvested culms are transformed into durable products. The increased lifespan of durable bamboo products made possible by modern technology can ensure that the sequestered carbon will not return quickly into the atmosphere, thereby prolonging the carbon storage of bamboo. According to Agarwal and his colleague [30] research result in the Mid Himalayan region conducting in comparing the carbon sequestration potential under different bamboo species, Monopodial species (*P. nigra*) has shown a high potential to sequester carbon than other Sympodial species. This might be due to high density of culms and high percent dry matter in the Himalayan region [31]. Currently, in China, about 53 M ha of forest plantation is there with a volume stock of 1.5 billion m³. Between 2005 and 2020, China has pledged to establish more than 40 million ha of plantations, referred to as carbon sink forest [32]. As plantations have been recognised as the national strategy for mitigating atmospheric CO₂, it is essential to assess the potential of fast-growing and high yield plantations in carbon storage and sequestration at the stand, regional and national scales (Fig. 1).

The area coverage of bamboo forests in China is about 6Mha, which stores about 780 Tg carbon, accounting for 14% of total forest carbon stock in China. Besides the carbon density of bamboo forest ecosystems in China, the estimated global bamboo carbon stock is about 4 Pg, accounting

for 0.43%-0.61% of total global forest carbon stock [33].

To combat climate change, bamboo should be a core development resource – providing

countries and development partners with a wealth of practical solutions to reduce the negative effects that changing climate patterns have on millions of rural communities (Fig. 2).




Fig. 1. Contribution of bamboo on adding C sink and reducing C emission (Source: Yuen [33])




Fig. 2. The five key functions of bamboo help to mitigate/adapt the impacts of climate change

6. SUMMARY

Currently, the concern of climate change is a severe and burning issue of global agendas. In this paper, we tried to review the contribution of bamboo forests to mitigating the impacts of climate change and the versatility of its ecological and socioeconomic development benefits. It offers one of the quickest ways to remove huge amounts of CO₂ from the atmosphere. It minimises CO₂ gas and generates more amount of oxygen than an equivalent stand of other tree species. Many scholars suggested that bamboo forest ecosystems provide significant services for human adaptation and development at the same time mitigating climate change impacts through carbon sequestration. Under well-managed bamboo plantations, it shows an effective carbon sink and better performance than other tree species growing under similar conditions. Despite this, it is a source of income in rewarding the diverse requirements at small and large-scales in rural areas. Generally, this review prepared to demonstrate the role of bamboo forests towards mitigating and adapting potential to overcome the impacts of climate change seen in the world and particularly to China. Therefore, advancing bamboo farming systems at different levels, it's advantageous in reducing greenhouse gas in the atmosphere and expanding bamboo forests in future under wider use and intensive management is recommended.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAO. Global forest resources assessment 2010: Main report. Food and agriculture Organization of the United Nations, Rome; 2010.
2. Janzen DH. Why bamboos wait so long to flower Annual? Rev. Ecol. Syst. 1976;7(1): 347–391. DOI:10. 1146/annurev.es. 07.110176. 002023.
3. Isagi Y, Kawahara T, Kamo K, Ito H. Net production and carbon cycling in a bamboo *Phyllostachys pubescens* stand. Plant Ecol. 1997;130:41–52. DOI:10.1023/A:1009711814070.
4. Li R, Werger MJA, During HJ, Zhong ZC. Carbon and nutrient dynamics in relation to growth rhythm in the giant bamboo *Phyllostachys pubescens*. Plant Soil. 1998;201(1):113–123. DOI:10.1023/A:1004322812651.
5. Lobovikov M, Lou YP, et al. The poor man's carbon sink. Bamboo in climate change and poverty alleviation. Non-Wood Forest Products. Working Document (FAO), no 8, FAO, Rome (Italy), Forestry Dept. 68p., FAO; 2009.
6. Zhu ZH. The development of bamboo and rattan in tropical China. Beijing, China: China Forestry Publishing House; 2001.
7. SFAPRC. Statistics of forest resources in China (1999–2003). State Forestry Administration, P. R. China; 2006.
8. IPCC. Solomon S, Qin D, Manning M. Chen Z, Marquis M, Averyt KBM. Tignor and Miller HL (eds). The Physical science basis, contribution of working group I to the fourth assessment report of the Intergovernmental Panel on Climate Change, Cambridge and New York, NY, USA. 2007;996.
9. Seppala R, Buck A, Katila. Eds. Adaptation of forests and people to climate change: A global assessment report. IUFRO World Series. Vienna, IUFRO. 2009;22.
10. Wei SJL, Sun SW, Wei, et al. Effects of climate changes on forest disasters and the preventive measures. Journal of Catastrophology. 2013;28(1):36-38.
11. MoEF (India. Ministry of Environment and Forests). India Second National Communication to the United Nations Framework Convention on Climate Change. New Delhi, Ministry of Environment and Forests, Government of India; 2012.
12. Hal Stern. Reviewed work(s): Source: The American Statistician. 2006;60(4):328-331.
13. Eliasch J. Climate change: Financing global forests: The Eliasch Review; 2008.
14. Dransfield S. Widjaja E. Eds. Bamboo plant resources of South-East Asia. No. 7. Leiden, Backhuys Publishers. 1995;190.
15. INBAR. Bamboo: A strategic resource for countries to reduce the effects of climate change. INBAR Policy Synthesis Report; 2009.
16. Magel E, Kruse S, Lutje G, Liese W. Soluble carbohydrates and acid lyertases involved in the rapid growth of the developing culms in *Sasa palmate* (Bean) Camus. Bamboo Science and Culture, Baton Rouge/USA. 2005;19(1)S.23-29.

17. Zhou GM, Jiang PK, Mo LF. Bamboo: A possible approach to the control of global warming. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009; 10(5):547-550.
18. Liese W. Bamboos as carbon sink fact or fiction? VIII World Bamboo Congress Proceedings. 2009;3:71-77.
19. FSI (Forestry Survey of India). India state of forestry report 2011. Dehradun, Forest Survey of India. 2011;286.
20. ICFRE (India Council of Forestry, Research and Education). Bamboo. Dehradun, ICFRE; 1992.
21. Banik RL. Issues of production of bamboo planting materials-Lessons and strategies. *Indian For (Bamb Iss)*. 2008;34(3):291-304.
22. Maoyi Fu, Banik RL. Bamboo, people and the management. Proceedings of the Vth International Bamboo Workshop and the IV International Bamboo Congress Ubud, Bali, Indonesia 10-22 June 1995. Volume I Propagation and management INBAR Technical Report No. 1995;8:18-30.
23. Terefe R, Samuel D, Sanbeto M, Daba M. Adaptation and growth performance of different lowland bamboo species in Bako, West Shoa, Ethiopia. *Journal of Natural Sciences Research*. 2016;6(9):61-65.
24. Lou Y, Li Y, Kathler B, Giles H, Zhou G. Bamboo and climate change mitigation: Comparative analysis of carbon sequestration. INBAR. Technical Report No. 32. 2010;1-20.
25. Isagi Y. Carbon stock and cycling in a bamboo *Phyllostachys bambusoides* stand. *Ecol. Res*. 1994;9:47-55.
26. Nath AJ, Lai R, Das AK. Above ground standing biomass for carbon storage in village bamboos in northeast India. *Bio. Bioeng*. 2009;33:1188-1196.
27. Janssen JJA. Building codes for bamboo housing. In Ganapathy M, Janssen JA, Sastry CB ed., *Bamboo, people and the environment. Engineering and utilization. Proceedings of the Vth International Bamboo Workshop*. 1996;3.
28. Kuehl Y, Yiping L. Carbon off- setting with bamboo. INBAR Working paper No. 71. Beijing, INBAR; 2012.
29. Zhou GM, Jiang PK. Density, storage and spatial distribution of carbon in *Phyllostachys pubescens* forest. *Sci. Sil. Sin*. 2004;40:20-25.
30. Agarwal A, Purwar JP. Evaluation of above ground biomass produced by *Dendrocalamus asper* in North Western Himalayan region of India. In: Proceedings of VIII World Bamboo Congress. 2009;4: 91-96.
31. Agarwal A, Purwar JP. Biomass production and carbon sequestration potential of various bamboo species in the Mid Himalayan Region of India. Paper Presented at World Bamboo Congress; 2012.
32. Chen X, Zhang X, Zhang Y, Booth T, He X. Changes of carbon stocks in bamboo stands in China during 100 years. *Forest Ecology & Management*. 2009;258:1489-1496.
33. Yuen JQ, Fung T, Ziegler AD. Carbon stocks in bamboo ecosystems worldwide: estimates and uncertainties. *For Ecol Management*. 2017;393:113-138.

© 2019 Terefe et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/44239>