

## Effect of Powdered and Composted Meat Bones on the Growth and Yield of Water Spinach (*Ipomoea aquatica*)

Badhon Ahmmed<sup>1\*</sup>, Tamzid Bin Shafique<sup>1</sup> and Shaikh Motasim Billah<sup>1</sup>

<sup>1</sup>Soil, Water and Environment Discipline, Khulna University, Khulna-9208, Bangladesh.

### Authors' contributions

This work was carried out in collaboration among all authors. Author BA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors TBS and SMB managed the analyses of the study. Authors BA and SMB managed the literature searches. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJSSPN/2019/v4i430052

#### Editor(s):

(1) Dr. Pankaj Gupta, Professor, Dolphin (PG) College of Science & Agriculture, Punjabi University, India.

#### Reviewers:

- (1) Jamile Da Silva Oliveira, Brazilian Agricultural Research Corporation, Brazil.  
(2) Rahul Datta, Mendel University, Czechia.  
(3) Jaime Cuauhtemoc Negrete, Autonomous Agrarian Antonio Narro University, Mexico.  
(4) Rebecca Yegon, University of Embu, Kenya.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50197>

Original Research Article

Received 05 May 2019  
Accepted 15 July 2019  
Published 01 August 2019

### ABSTRACT

For agriculture, the most important is the soil's function as a base and medium for plant growth. The soil health and soil condition are important for plant growth. And the management of soil organic matter is an important part in case of managing soil health and maintaining soil conditions. The experiment was carried out in the field lab (Net House) of Soil, Water and Environment discipline, Khulna University, Bangladesh. Total 21 plastic pots were used to continue the experiment with 7 treatments. For the experiment, the meat bones were collected and used in the experimental pot along with soil in different doses. This research has revealed that meat bones both powdered and composted, has significantly increased (about 20-25%) the growth parameters of water spinach (*Ipomoea aquatica*) such as a number of leaves, shoot length, fresh weight, dry weight and moisture content. So, the application of powdered and composted meat bones in soil will contribute in the growth of water spinach (*Ipomoea aquatica*).

\*Corresponding author: Email: badhonahmmed1996@gmail.com;

**Keywords:** Soil organic matter; meat bones; water spinach (*Ipomoea aquatica*); pot experiment.

## 1. INTRODUCTION

Agriculture is the most important sector of the economy of Bangladesh, contributing about 23% of the country's GDP and employing about 62% of the total labour force. Bangladesh, after independence, has adopted various measures to raise vegetable production for tackling the malnutrition problem in the country.

Water spinach (*Ipomoea aquatica*) also known as 'kalmi shak' (in Bengali) is one of the popular vegetables that is promoted to grow in Bangladesh due to its good nutritive value, antioxidant properties, high fiber content and many other health-related benefits [1]. It is a vascular semi-aquatic herbaceous perennial plant belonging to *Convolvulaceae* [2]. Water spinach, *Ipomoea aquatica* Forssk. (*Convolvulaceae*), is an aquatic or semi-aquatic edible herb [3]. It has a hollow and viny stem, grows prostrate or floating, and roots are coming from the nodes that penetrate the soil. Water spinach is native to the tropics and subtropics of Southeast Asia, Southern China and India [4], [5]. It is a green vegetable and is ranked high among the world's healthiest foods, and there are plenty of reasons for it being so. It is one of the ideal options to manage weight and to lower the cholesterol levels. It consists of rich amounts of iron that is required by the red blood cells during process of hemoglobin formation. So, people who are suffering with anemia should include their diet with this iron rich leafy vegetable [6].

For agriculture, the most important is the soil's function as a base and medium for plant growth. Soil health and soil condition is important, and the management of soil organic matter is an important part of managing soil health and maintaining soil condition. Most of the soils of Bangladesh have low organic matter content, usually less than 2% [7]. A good soil should have at least 2.5% organic matter, but in Bangladesh, most soils have less than 1.5%, and some soils have even less than 1% organic matter [8]. The pressure on the management of soil organic matter is increasing as costs of inputs for agriculture increase and the capacity and ability to overcome soils in poor condition by adding more fertilizer, adding one more cultivation, adding one more irrigation or adding another input are diminished [9]. The ultimate source of organic matter for most soils is through the fixation of carbon dioxide from the atmosphere through photosynthetic reactions by plants. There is also a very small input from autotrophic bacteria. However, in some instances, there may also be some input from industrial and mining products derived from petroleum or coal. At the broad scale, these sources of soil carbon are insignificant. Soil organic matter is derived from organic materials that are added to the soil, and the majority of soil organic matter derives from the breakdown of residues remaining after plants have died. These residues can take the form of root residues located in the soil matrix or leaves, stems and stubble existing as litter on the soil surface. Animals also provide a proportion of the soil organic matter to varying degrees depending on management and the ecosystem [10].



**Fig. 1. Water spinach**

Additions of large amounts of organic materials as composts or as biochar can increase the levels of soil organic matter in soils [11]. Soils are used as a filter or sink for effluent and waste materials. The wastes produced from meat bones can be used as such types of organic materials as the replacement of compost or fertilizers.

Maximum people eat various types of meat daily. There are some particular occasions such as wedding ceremony, some religious festivals like Eid, Puja etc which is incomplete without meat. A lot of meat bones can be found after these ceremonies. If we use these bones as fertilizers for plant growth, then the waste load will be controlled. Usually, beef bones, however, other types of a creature like chicken or pigeon bone meal or emulsion can be used to prepare meat bone fertilizer. The processed bone meal goes through to become a powder or compost will quickly kill off many pathogens.

Meat and bone meal (MBM) can be a viable alternative to mineral fertilizers because it contains about 8% Nitrogen (N), 5 % Phosphorus (P), 1 % Potassium (K) and 10 % Calcium (Ca) [12], which makes it a valuable source of nutrients for plant production. It has about 50% protein, 35% ash, 8-12% fat, and 4-7% moisture, and contains a big amount of nutrients. Chemical properties of Meat Bone Materials (MBM) vary a lot from different raw materials. On average, the pH tends to be acidic, about 6.5. Organic matter in content is about 50% [13]. Besides a great deal of phosphorous, bone meal also provides smaller amounts of essential nitrogen, potassium and calcium. As the bone meal breaks down over the season, it will steadily and slowly release these nutrients for plant uptake. The advantageous aspects of slow release nutrition are that plant roots will not burn, and that soil vitality is increased for extended periods [14].

### 1.1 Objectives

To minimize the rate of waste load by using the meat bones as organic fertilizer which is environmental friendly as well as to improve soil health, for determining the effectiveness of different types of meat bones in water spinach cultivation, to assess the effect of composted and powdered meat bones on the growth and yield of water spinach (*Ipomoea aquatica*).

## 2. MATERIALS AND METHODS

### 2.1 Location of the Study Area

The experiment was carried out in the field lab (Net House) of Soil, Water and Environment discipline, Khulna University, Bangladesh.

### 2.2 Collection and Preparation of the Soil

The soil for our research was collected from the research field of Soil, Water and Environment Discipline, Khulna University. Spade, plastic bag was used for collecting the soil. Then the soil was air dried at room temperature, and all the plant debris were removed manually. Then the massive aggregates were broken by gentle crushing by a hammer. Then the soil was sieved by 2 mm sieve.

### 2.3 Preparation of Pot

After sieving, the soil was taken into pots for sowing the seeds. Total 21 plastic pots were used to continue the experiment. The pots were cleaned and labelled. Each Pot was filled with 3kg sieved soil. Pots had no pore in its bottom to protect the leaching of meat bone fertilizers from the soil. The experiment was started on 24<sup>th</sup> June, 2018.

### 2.4 Powdered and Composted Meat Bones Preparation

The meat bones were collected after eating. Then some of the bones were crushed into powder form with the help of crusher (Haman dista). To prepare meat bone compost, bones (pigeon, chicken, and beef) were crushed and transferred it into a plastic bag. Sufficient amount of water was added into it to degrade the meat bones and kept it by digging soil into 30 cm depth. After 60 days (4<sup>th</sup> June – 4<sup>th</sup> August), the compost was ready to be applied into my experiment. 10 g powdered and composted meat bones were applied to each pot. 5 g was mixed up with the soil before sowing the seeds, and the rest 5 g was added when the seeds were grown up to plants.

### 2.5 Treatments of Investigation

Seven treatments were used in the experiment.

T<sub>0</sub> = Control  
T<sub>1</sub> = 500 kg ha<sup>-1</sup> powdered beef bone

T<sub>2</sub> = 500 kg ha<sup>-1</sup> powdered chicken bone  
T<sub>3</sub> = 500 kg ha<sup>-1</sup> powdered pigeon bone  
T<sub>4</sub> = 500 kg ha<sup>-1</sup> composted beef bone  
T<sub>5</sub> = 500 kg ha<sup>-1</sup> composted chicken bone  
T<sub>6</sub> = 500 kg ha<sup>-1</sup> composted pigeon bone

tagged separately, weighted, oven dried at 65°C temperature for 24 hours until moisture content reached to a minimum condition. The dried material of plants per pot from each treatment was collected.

## 2.6 Sowing of the Seeds

The seeds were sown on 24<sup>th</sup> June, 2018. The seeds were sown thoroughly as it was possible to keep uniformity and then the seeds were covered by soils. 0.01 g seeds (5 kg ha<sup>-1</sup> as recommended by BARI, 2005 for trial experiment) were sown in each pot and maximum seeds germinated within 5 days. After germination only five plants were kept in each pot.

### 2.7.3.1 Number of leaves

The number of leaves of five plants of each pot was counted and average value was taken.

### 2.7.3.2 Shoot length (cm)

Shoot length was measured using a measuring scale from the root level to the tip of the plant. From each pot, five plants were measured and averaged.

## 2.7 Intercultural Operations

### 2.7.1 Watering

For normal plant growth, the water was added regularly (2-3 days intervals) to each pot for maintaining enough moisture.

### 2.7.3.3 Root length (cm)

Root length was measured using a measuring scale from root level to the tip of the longest root at harvest, and their average value was taken as the root length in cm.

### 2.7.2 General observation

The pots under experiment were frequently observed to note any change in the crop growth and other characteristics. The crop growth was very satisfactory in some treatments. But some treatments showed a lower number of plants.

### 2.7.3.4 Fresh weight per plant (gm)

Harvest of five plants from each pot, fresh weight of the whole plant was taken by an electrical balance, and their mean value was calculated as fresh weight expressed in gm/plant.

### 2.7.3 Harvesting

The experimental crops were harvested after 40 days of germination. The harvested plants were

### 2.7.3.5 Dry weight per plant (gm)

Five plants from each pot were collected and oven dried at 65°C for 48 hours, weighed in gm/plant by an electrical balance and average value was recorded.



Fig. 2. Sowing the seeds of water spinach

### 2.7.3.6 Moisture content (%)

Percent moisture was calculated by using the formula:

$$\text{Moisture content (\%)} = \frac{W_f - W_o}{W_f} \times 100$$

Where,

$W_f$  = Fresh weight of the plant sample  
 $W_o$  = Oven dry weight of the plant sample

## 2.8 Analysis

The collected data on different parameters were represented in bar diagram by Microsoft office excel program 16.0 and statistically analyzed following the analysis of variance (ANOVA) technique. The Completely Randomized Design (CRD) was followed as experimental design and the different doses of meat bones in composted and powdered forms were the treatments.

## 3. RESULTS AND DISCUSSION

Effect of powdered and composted meat bone on the growth and yield of Water spinach (*Ipomoea aquatica*) was studied following pot experiment. The number of leaves per plant, shoot length per plant, root length per plant, fresh weight per plant, dry weight per plant and percent moisture content were measured for the plants treated with 10g powdered and composted meat bone and compared with plants grown with no meat bone which was control experiment. The data of morphological attributes are presented in Table 1.

### 3.1 Number of Leaves per Plant

The number of leaves of water spinach was significantly influenced by different treatments in maximum cases. The highest number of leaves was found in  $T_5$  (10.73) (Table 1). The results exhibited that there was the insignificant difference in  $T_1$ ,  $T_2$ ,  $T_4$  and  $T_6$  compared to the control experiment  $T_0$  and a significant difference was found in  $T_3$  and  $T_5$  (Fig. 3). Application of composted chicken bone showed a significant increment of the number of leaves in water spinach, whereas application of other meat bone fertilizers did not show any positive effect on the number of leaves of water spinach. On the basis of the number of leaves of plants, application of composted chicken bone is the best for using,

and it is recommended to use as a soil amendment for the growth of such type of leafy vegetables like water spinach.

### 3.2 Root Length per Plant (cm)

Length of root per plant ranged from 8.81cm to 13.91 cm and highest length of root found in  $T_6$  (13.91 cm) (Table 1). Length of root varied significantly in  $T_6$  compared with control treatment  $T_0$  and there was the insignificant difference among  $T_1$ ,  $T_2$ ,  $T_3$ ,  $T_4$  and  $T_5$  (Fig. 4). A distinct difference was found between  $T_0$  and  $T_6$ . So, application of compost of pigeon bone showed a significant increment of length of root in water spinach whereas application of powdered chicken bone, beef bones did not show any positive effect on the root length of water spinach. On the basis of the length of the root of plants application of composted pigeon bone is the best for uses, and it can be recommended for the plants.

### 3.3 Shoot Length per Plant (cm)

Length of shoot per plant ranged from 17.64 cm to 24.96 cm. and highest length of shoot was found in  $T_5$  (24.96 cm) (Table 1). There was insignificant relationship between  $T_1$ ,  $T_2$  and  $T_6$  but significant difference was found in  $T_5$  compared with the control treatment  $T_0$  (Fig. 5). So, application of composted chicken bone showed significant increment of length of shoot in water spinach whereas application of other meat bone fertilizers did not show any positive effect on the shoot length of water spinach. On the basis of the length of shoot of plants application of composted chicken bone is the best for uses and it is recommended for the plants.


### 3.4 Fresh Weight per Plant (g)

Fresh weight per plant ranged from 1.09 g to 3.47 g and highest fresh weight of plant was found in  $T_5$  (3.47 g) (Table 1). Fresh weight varied significantly in  $T_4$ ,  $T_5$ ,  $T_6$  compared with control treatment  $T_0$  but there was the insignificant difference between  $T_0$ ,  $T_1$ ,  $T_2$  and  $T_3$  (Fig. 6) The observation was  $T_5 > T_6 > T_4 > T_1 > T_2 > T_3 > T_0$ . So, application of compost of chicken bones showed a significant increment of fresh weight in water spinach, whereas application of meat bones and pigeon bones did not show any positive effect on the fresh weight. On the basis of the fresh weight of plants application of composted chicken bone is the best for uses, and it is recommended for the leafy vegetables.




**Table 1. Observed growth factors of the experimented water spinach for different treatments**

Treatments	No. of leaves /plant	Root length /plant (cm)	Shoot length /plant (cm)	Fresh weight /plant (g)	Dry weight /plant (g)	Moisture content (%)
T <sub>0</sub>	8	8.81	17.64	1.09	0.08	64.36
T <sub>1</sub>	9	9.31	24.32	1.56	0.37	76.49
T <sub>2</sub>	9	9.98	24.25	1.23	0.32	73.85
T <sub>3</sub>	8	10.93	18.65	1.17	0.35	70.25
T <sub>4</sub>	9	12.65	22.01	2.67	0.40	84.97
T <sub>5</sub>	11	13.51	24.96	3.47	0.52	85.05
T <sub>6</sub>	10	13.91	23.93	3.30	0.48	85.39



**Fig. 3. Effect of powdered and composted meat bone on leaves number of water spinach**



**Fig. 4. Effect of powdered and composted meat bone on root length of water spinach**




Fig. 5. Effect of powdered and composted meat bone on shoot length of water spinach




Fig. 6. Effect of powdered and composted meat bone on the fresh weight of water spinach

### 3.5 Dry Weight per Plant (g)

Dry weight per plant ranged from 0.08g to 0.52g and highest dry weight was found in T<sub>5</sub> (0.52g) (Table 1). Dry weight varied significantly in all treatments compared to the control treatment T<sub>0</sub> and there was an insignificant difference between T<sub>2</sub> and T<sub>3</sub> as well as T<sub>1</sub> and T<sub>4</sub> (Fig. 7). So, application of compost of chicken bone showed significant increment of the dry weight in water spinach whereas application of other meat

bone fertilizers did not show any positive effect on the dry weight. On the basis of dry weight of plants application of composted chicken bone is the best for using and it is recommended for the plants.

### 3.6 Moisture Content per Plant (%)

Moisture content per plant ranged from 64.36% to 85.39% and highest moisture content found in T<sub>6</sub> (85.39%) and the lowest moisture content

observed in T<sub>0</sub> (Table 1). The observations are T<sub>6</sub> > T<sub>5</sub> > T<sub>4</sub> > T<sub>1</sub> > T<sub>2</sub> > T<sub>3</sub> > T<sub>0</sub>. The following figure shows significant differences when control experiment is compared with other treatment and significant differences are found in T<sub>1</sub> and T<sub>3</sub> but insignificant relationship was found among T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub> treatments (Fig. 8). So, application of compost of pigeon bone showed a significant

increment of moisture content of water spinach whereas application of chicken bone, beef bones both powdered and composted did not show any positive effect on the moisture content of water spinach. On the basis of the moisture content of plants application of composted pigeon bone is the best for using and it can be recommended for the plants.




Fig. 7. Effect of powdered and composted meat bone on the dry weight of water spinach




Fig. 8. Effect of powdered and composted meat bone on the moisture content of water spinach



#### 4. SUMMARY AND CONCLUSION

Powdered and composted meat bones contain sufficient nutrients such as nitrogen, phosphorus, potassium, calcium which are capable to enhance the growth rate of plant. This research has revealed that meat bones both powdered and composted, has significantly increased the growth parameters of water spinach (*Ipomoea aquatica*) such as number of leaves, shoot length, fresh weight, dry weight and moisture content. The result showed that these parameters are mostly increased by using composted chicken bones compared to that of other treatments. Only root length and moisture content of water spinach is increased by using composted pigeon bones. The highest vegetative growth was found by using composted chicken bones, and lowest vegetative growth was found at control experiment. So, the use of composted chicken bone has the potential to show the highest response on the growth and yield of water spinach and other types of leafy vegetables.

So, it is recommended for the farmers to use composted meat bones specially composted chicken bones for the improvement of soil quality and to improve its productivity. On the contrary, a load of waste from meat bones will be reduced and will help in waste management. In the same time, the use of costly and harmful chemical fertilizers will be reduced and will introduce organic farming, which is environment friendly and beneficial for human health.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Hongfei F. Evaluation of antioxidant activities of principal carotenoids available in water spinach (*Ipomoea aquatica*). Journal of Food Composition and Analysis. 2011;24(2):288-297.
2. USDA (The United States Department of Agriculture), *Ipomoea aquatica* Forssk; 2005.
3. Dua TK, Dewanjee S, Gangopadhyay M, Khanra R, Zia-Ul-Haq M, DeFeo V. Ameliorative effect of water spinach, *Ipomoea aquatic* (Convolvulaceae), against experimentally induced arsenic toxicity. Journal of Translational Medicine. 2015;13:81.  
DOI: 10.1186/s12967-015-0430-3
4. Gothberg A, Greger M, Bengtsson B. Accumulation of heavy metals in water Spinach (*Ipomoea aquatica*) cultivated in Bangkok region, Thailand. Environmental Toxicology and Chemistry. 2005;21:1934-1939.
5. Chen BH, Yang SH, Han H. Characterization of major carotenoids in water Convolvulus (*Ipomoea aquatica*) by open-column, thin layer and high-performance Liquid chromatography. Journal of Chromatography. 1991;534: 147-155.
6. Gupta S, Lakshmi AJ, Manjunath MN, Prakash J. Analysis of nutrient and antinutrient content of underutilized green leafy vegetables. LWT Food Science. 2005;43:11-21.
7. Bhuiyan NI. Crop production trend and need of sustainability in agriculture. A paper presented in a three-day long workshop on integrated nutrients management for sustainable agriculture held at SRDI, Dhaka, June, 26-28; 1994.
8. BARC (Bangladesh Agricultural Research Council). Fertilizer Recommendation Guide. 2005;22, 48, 174.
9. MacEwan RJ. Soil Health for Victoria's Agriculture- Context, Terminology and Concepts. MIS 07898 Final Report. Department of Primary Industries Research Victoria Bendigo, State of Victoria; 2007.
10. Tate, Robert L. Soil organic matter – biological and ecological effects. John Wiley and Sons, New York; 1987.
11. Gibson TS, Chan KY, Sharma G, Shearman R. Soil Carbon Sequestration Utilizing Recycled Organics - A review of the scientific literature. Project 00/01R-3.2.6A. The Organic Waste Recycling Unit, NSW Agriculture. Report prepared for Resource NSW; 2002.
12. Ylivainio K, Uusitalo R, Turtola E. Meat bone meal and fox manure as P sources for ryegrass (*Lolium multiflorum*) grown on a limed soil. Nutrient Cycling in Agroecosystems. 2007;81:267-278.

13. Jeng AS, Haraldsen TK, Gronlund A, Pedersen PA. Meat and bone meal as nitrogen and phosphorus fertilizer to cereals and rye grass. *Nutrient Cycling in Agroecosystems*. 2006;76:183-191.
14. Cayuela ML, Sinicco T, Mondini C. Mineralization dynamic and biochemical properties during initial decomposition of plant and animal residues in soil. *Applied Soil Ecology Journal*. 2009;41:118–127.

---

© 2019 Ahmed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/50197>